

正誤表

『ニューロンの生理学』に下記の誤りがありました。
謹んでお詫び申し上げますとともに、訂正いたします。

●492 ページ 14 行目、497 ページ見出し

誤 15.3 G タンパク共役型または代謝型グルタミン酸受容体

正 15.3 グルタミン酸を介するシナプス後部での脱分極：異なるタイプの受容体の役割

●参考文献の脱落

・158 ページ (付録 4.1)

参考文献

[A4.1.1] Caine SB. Cocaine abuse : hard knocks for the dopamine hypothesis? Nature Neurosciences 1998 ; 1 , 90-2.

[A4.1.2] Giros B, Jaber M, Jones SR, Wightman RM, Caron MG. Hyperlocomotion and indifference to cocaine and amphetamine in mice lacking the dopamine transporter. Nature 1996 ; 379 : 606-12.

[A4.1.3] Koob GF. Drugs of abuse : anatomy, pharmacology and function of reward pathways. TIPS 1992 ; 13 : 177-84

[A4.1.4] Sokoloff P, Schwartz JC. Novel dopamine receptors half a decade later. TIPS 1995 ; 16 : 270-5.

[A4.1.5] Stip E, Godbout R. Cortex préfrontal, monoamines et neuropsychologie de la schizophrénie. Med Sci 1995 ; 11 : 1531-9.

[A4.1.6] Woolverton WL, Jonhson KM. Neurobiology of cocaine abuse. TIPS 1992 ; 13 : 193-200.

・192 ページ (付録 5.1)

参考文献

[A5.1.1] Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. Pflügers Arch. 1981 ; 391 : 85-100.

・196 ページ (付録 5.2)

参考文献

[A5.2.1] Toth PT, Miller RJ. Calcium and sodium currents evoked by action potential waveforms in rats sympathetic neurons. *J Physiol* 1995 ; 485 : 43-57.

[A8.1.1] Akopian AN, Sivilotti L, Wood JN. A tetrodotoxin-resistant voltage-gated sodium channel expressed by sensory neurons. *Nature* 1996 ; 379 : 257-62.

・ 280 ページ (付録 8. 1)

参考文献

[A8.1.2] Alonso A, Llinas RR. Subthreshold Na⁺ -dependent theta-like rhythmicity in stellate cells of entorhinal cortex layer II. *Nature* 1989 ; 342 : 175-7.

[A8.1.3] Catterall WA. Structure and function of voltage-gated ion channels. *Ann Rev Biochem* 1995 ; 64 : 493-531.

[A8.1.4] Chao TI, Alzheimer C. Effects of phenytoin on the persistent Na⁺ current of mammalian CNS neurones. *Neuroreport* 1995 ; 6 : 1778-80.

[A8.1.5] Crill WE. Persistent sodium current in mammalian central neurons. *Ann Rev Physiol* 1996 ; 58 : 349-62.

[A8.1.6] Dubois JM, Bergman C. Late sodium current in the node of Ranvier. *Eur J Physiol* 1975 ; 357 : 145-8.

[A8.1.7] Hoehn K, Watson TWJ, Mac Vicar BA. A novel tetrodotoxin-insensitive, slow sodium current in striatal and hippocampal neurones. *Neuron* 1993 ; 10 : 543-52.

[A8.1.8] Kostyuk PG, Veselovsky NS, Tsyndrenko AY. Ionic currents in the somatic membrane of rat dorsal root ganglion neurons. I. Sodium Currents. *Neurosci* 1981 ; 6 : 2423-30.

[A8.1.9] Oka Y. Tetrodotoxin-resistant persistent Na⁺ current underlying pacemaker potentials of fish gonadotrophin-releasing hormone neurones. *J Physiol* 1995 ; 482 : 1-6.

[A8.1.10] Oka Y. Characterization of TTX-resistant persistent Na⁺ current underlying pacemaker potentials of fish gonadotrophin-releasing hormone (GnRH) neurons. *J Neurophysiol* 1996 ; 75 : 2397-404.

[A8.1.11] Rogart RB, Cribbs LL, Muglia LK, Dephart DD, Kaiser MW. Molecular cloning of a putative tetrodotoxin-resistant rat heart Na⁺ channel isoform. *PNAS*

1989 ; 86 : 8170-4.

[A8.1.12] Roy ML, Narahashi T. Differential properties of tetrodotoxin-sensitive and tetrodotoxin-resistant sodium channels in rat dorsal root ganglion neurons. *J Neurosci* 1992 ; 12 : 2104-11.

[A8.1.13] Taylor CP. **Na⁺ currents that fail to inactivate.** *TINS* 1993 ; 16 : 455-60.

[A8.1.14] Yoshida S. **Tetrodotoxin-resistant sodium channels.** *Cell Mol Neurobiol* 1994 ; 14 : 227-44.

[A8.1.15] Yoshida S, Matsuda Y, Samejima A. Tetrodotoxin-resistant sodium and calcium components of action potentials in dorsal root ganglion cells of the adult mouse. *J Neurophysiol* 1978 ; 41 : 1096-106.

・ 305 ページ(第 9 章)

参考文献

[9.1] Amoroso S, Schmid-Antomarchi H, Fosset M, Lazdunski M. Glucose, sulfonylureas, and neurotransmitter release : role of ATP-sensitive potassium channels. *Science* 1990 ; 247 : 852-4.

[9.2] Chesnoy-Marchais D. A Cl⁻ conductance activated by hyperpolarization in *Aplysia* neurones. *Nature* 1982 ; 299 : 359-61 .

[9.3] Chesnoy-Marchais D. Characterization of a chloride conductance activated by hyperpolarization in *Aplysia* neurones. *J Physiol* 1983 ; 342 : 277-308.

[9.4] Chesnoy-Marchais D, Evans M. Chloride channels activated by hyperpolarization in *Aplysia* neurones. *Eur J Physiol* 1986 ; 407 : 694-6.

[9.5] Chesnoy-Marchais D, Fritsch J. Activation by hyperpolarization and atypical osmosensitivity of a Cl⁻current in rat osteoblastic cells. *J Memb Biol* 1994 ; 140 : 173-88.

[9.6] Di Francesco D. **Pacemaker mechanisms in cardiac tissue.** *Ann Rev Physiol* 1993 ; 55 : 455-72.

[9.7] Di Francesco D, Ferroni A, Mazzanti M, Tromba C. Properties of the hyperpolarizing-activated current (*i_p*) in cells isolated from the rabbit sino-atrial node. *J Physiol* 1986 ; 377 : 61-88.

[9.8] Di Francesco D, Mangoni M. Modulation of single hyperpolarization-activated channels (i_f) by AMP cyclique in the rabbit sino-atrial node. *J Physiol* 1994 ; 474 : 473-82.

[9.9] Doupnik CA, Davidson N, Lester HA. The inward rectifier potassium channel family. *Curr Opin Neurobiol* 1995 ; 5 : 268-77.

[9.10] Gauss R, Seifert R, Kaupp B. Molecular identification of a hyperpolarization-activated channel in sea urchin sperm. *Nature* 1998 ; 393 : 583-7.

[9.11] Hagiwara S, Miyazaki S, Rosenthal N. Potassium current and the effect of cesium on this current during anomalous rectification of the egg cell membrane of a starfish. *J Gen Physiol* 1976 ; 67 : 621-38.

[9.12] Katz B. Les constantes électriques de la membrane du muscle. *Arch Sci Physiol* 1949 ; 3 : 285-300.

[9.13] Kofuji P, Doupnik CA, Davidson N, Lester HA. A unique P-region residue is required for slow voltage-dependent gating of a G protein-activated inward rectifier potassium channel expressed in *Xenopus* oocytes. *J Physiol* 1996 ; 490 : 633-45.

[9.14] Lopatin AN, Makhina EN, Nichols CG. Potassium channel block by cytoplasmic polyamines as the mechanism of intrinsic rectification. *Nature* 1994 ; 372 : 366-9.

[9.15] Lu Z, MacKinnon R. Electrostatic tuning of Mg^{2+} affinity in an inward-rectifier potassium channel. *Nature* 1994 ; 371 : 243-6.

[9.16] Ludwig A, Zong X, Jeglitsch M, Hofmann F, Biel M. A family of hyperpolarization-activated mammalian cation channels. *Nature* 1998 ; 393 : 587-91.

[9.17] Mc Cormick DA, Pape HC. Properties of a hyperpolarization-activated cation current and its role in rhythmic oscillation in thalamic relay neurones. *J Physiol* 1990 ; 431 : 291-318.

[9.18] Miller C. Open state substructure of single chloride channels from *Torpedo* electroplax. *Phil Trans R Soc Lond. B* 1982 ; 299 : 401-11.

[9.19] Silver MR, De Coursey TE. Intrinsic gating of inward rectifier in bovine pulmonary artery endothelial cells in the presence or absence of internal Mg^{2+} . *J Gen Physiol* 1990 ; 96 : 109-33.

[9.20] Soltesz I, Lightowler S, Leresche N, Jassik-Gerschenfeld D, Pollard CE,

Crunelli V. Two inward currents and the transformation of low-frequency oscillations of rat and cat thalamo-cortical cells. *J Physiol* 1991 ; 441 : 175-97.

[9.21] Staley K. The role of an inwardly rectifying chloride conductance in postsynaptic inhibition. *J Neurophysiol* 1994 ; 72 : 273-84.

[9.22] Stanfield PR, Nakajima Y, Yamaguchi K. Substance P raises neuronal membrane excitability by reducing inward rectification. *Nature* 1985 ; 315 : 498-501.

[9.23] Surprenant A, North RA. Mechanism of synaptic inhibition by noradrenaline acting at α_2 -adrenoreceptors. *Proc R Soc Lond B* 1988 ; 234 : 85-114.

[9.24] Thieman A, Gründer S, Push M, Jentsch T. A chloride channel widely expressed in epithelial and non-epithelial cells. *Nature* 1992 ; 356 : 57-60.

[9.25] Vandenberg CA. Inward rectification of a potassium channel in cardiac ventricular cells depends on internal magnesium ions. *PNAS* 1987 ; 84 : 2560-4.

・ 308 ページ (付録 9.1) [305 ページより移動]
参考文献

[A9.1.1] Amoroso S, Schmid-Antomarchi H, Fosset M, Lazdunski M. Glucose, sulfonylureas, and neurotransmitter release : role of ATP-sensitive potassium channels. *Science* 1990 ; 247 : 852-4.

[A9.1.2] Ashcroft FM. Adenosine 5'-triphosphate-sensitive potassium channels. *Ann Rev Neurosci* 1988 ; 11 : 97-118.

[A9.1.3] Ashford MLJ, Sturgess NC, Trout NJ, Gardner NJ, Hales CN. Adenosine-5'-triphosphate-sensitive ion channels in neonatal rat cultured central neurones. *Eur J Physiol* 1988 ; 412 : 297-304.

[A9.1.4] Ashford MLJ, Boden PR, Treherne JM. Glucose-induced excitation of hypothalamic neurones is mediated by ATP-sensitive K⁺ channels. *Eur J Physiol* 1990 ; 415 : 479-83.

[A9.1.5] Edwards G, Weston AH. Pharmacology of the potassium channel openers. *Cardiovascular Drugs and Therapy* 1995 ; 9 : 185-93.

[A9.1.6] Inagaki N, Gono T, Clement IV JP, Namba N, Inazawa J, Gonzalez G *et al.*

Reconstitution of K_{ATP} : an inward rectifier subunit plus the sulfonylurea receptor.
Science 1995 ; 270 : 1166-70.